



Oddfellows Arms, 25 Main St, Higham on the Hill,
Nuneaton, CV13 6AE

Dusk Emergence Survey for Bats



July 2024

All Ecology Ltd

Tel: 01453 393001

Email: info@allecology.co.uk

Web: www.allecology.co.uk

© ALL ECOLOGY 2024

Document Control

Site: Oddfellows Arms, 25 Main St, Higham on the Hill, Nuneaton, CV13 6AE

Title: Dusk Emergence Survey for Bats

For: Cartwright Homes

Project Number: 1976

Document Version: 1.0

Survey Date(s): 11th July 2024

Document Date: 21st July 2024

Version	Date	Version Details	Prepared by	Reviewed by	Approved by
1.0	11/07/24	-	DR	JG	JG



England and Wales Office

All Ecology Ltd,
Haw Street,
Coaley,
Dursley,
Gloucestershire,
GL11 5AY.

Scotland Office

All Ecology Ltd,
Bowfield Road,
Howwood,
Renfrewshire,
PA9 1BZ.

Tel: 01453 393001
Email: info@allecology.co.uk
Web: www.allecology.co.uk

All Ecology is the trading name of All Ecology Ltd. Registered in England and Wales, Company Number 8306310

Notice to Readers:

The results of the survey and assessment work undertaken by All Ecology are representative at the time of surveying.

Every endeavour has been made to identify the presence of protected species on site, where this falls within the agreed scope of works.

The flora and fauna detailed within this report are those noted during the field survey and from anecdotal evidence. It should not be viewed as a complete list of flora and fauna species that may frequent or exist on site at other times of the year.

Up to date standard methodologies have been used, which are accepted by Natural England and other statutory conservation bodies. No responsibility will be accepted where these methodologies fail to identify all species on-site.

All Ecology cannot take responsibility where Government, national bodies or industry subsequently modify standards.

All Ecology cannot accept responsibility for data collected from third parties.

Reference to sections or particular paragraphs of this document taken out of context may lead to misrepresentation.

Contents

Contents	1
1.0 Introduction	3
Background	3
Objectives and Aim	3
Site Location	4
2.0 Legislation and Status	5
Bats	5
3.0 Methodology	7
Personnel	7
Dusk Emergence Survey	7
Equipment	7
Assessment	7
Limitations	9
4.0 Results	10
Dusk Emergence Survey	10
5.0 Evaluation	11
Dusk Emergence Survey	11
Site Status Assessment	11
6.0 Impacts and Recommendations	12
Impacts	12
Further Surveys	12
Legal Compliance	12
Mitigation	12
7.0 References	14

8.0 Plans

15

1.0 Introduction

Background

- 1.1 All Ecology was commissioned to undertake a Dusk Emergence Survey for Bats on a disused public house known as the Oddfellows Arms, 25 Main St, Higham on the Hill, Nuneaton, CV13 6AE.
- 1.2 The proposal to redevelop the site includes the conversion of the current building to provide two new dwellings.
- 1.3 A previous dusk emergence survey of the building was carried out by Guma in August 2019. This survey is now out of date and the present survey was requested to provide an up to date assessment.
- 1.4 In accordance with the Bat Surveys for Professional Ecologists Good Practice Guidelines 4th Edition (Collins, 2023), where a building is deemed to have low potential for roosting bats, the building should be subject to one dusk emergence survey between May – August inclusive to give confidence in a negative result.

Objectives and Aim

- 1.5 The main objectives and aim of the survey were to establish the following:
 - Presence/absence of bat roosts.
 - Status of roosts if present.
 - Whether a European Protected Species (EPS) licence is required to ensure legal compliance.
 - Which type of mitigation measures would need to be employed.

Site Location



Figure 1: Site location plan (buildings indicated).

2.0 Legislation and Status

Bats

- 2.1 All species of bat are listed on Schedule 5 of The Wildlife and Countryside Act (1981) and as such receive protection under Section 9 of this Act. This has been amended several times, most recently by the Countryside and Rights of Way Act 2000, which added 'or recklessly' to Section 9(4) (a) and (b). In summary, it is a criminal offence to:
- Intentionally kill, injure or take a wild bat.
 - Be in possession of, or control, any live or dead wild bat or part of, or anything derived from a wild bat.
 - Intentionally or recklessly damage, destroy or obstruct access to any place that a wild bat uses for shelter or protection.
 - Intentionally or recklessly disturb any wild bat whilst it is occupying a structure or place that it uses for shelter or protection.
 - Transport for sale or exchange, or offer for sale or exchange, a live or dead bat or any part of a bat.
- 2.2 The Conservation of Habitats and Species Regulations 2010, consolidate all the various amendments made to the Conservation (Natural Habitats, &c.) Regulations 1994, in respect of England and Wales. It is an offence to possess, sell or offer or transport for sale any European species of bat or any part derived from such a species. These Regulations also remove the 'incidental result defence'. In other words, it is no longer a defence to show that the killing, capture or disturbance of a species covered by the Regulations or the destruction or damage of their breeding sites or resting places was the incidental and unavoidable result of a lawful activity. Natural England can grant European Protected Species (EPS) licenses in respect of development to permit activities that would otherwise be unlawful.
- 2.3 Under Section 40 of the Natural Environment and Rural Communities Act (2006) public bodies, including Local and Regional Planning Authorities have a duty to 'have regard' to the conservation of biodiversity in England when carrying out their normal functions, which includes consideration of planning applications. In compliance with Section 41 of the Act, the Secretary of State has published a list of species considered to be of principal importance for conserving biodiversity in England. This is known as The England Biodiversity List, all of which make up the UK BAP Priority Species. This list forms the basis of the UK Biodiversity Framework, and in England, Biodiversity 2020: A strategy for England's wildlife and ecosystem services (Defra, 2011). Regional Planning Bodies and Local Planning Authorities will use it to identify the species that should be afforded priority when applying the requirements of the National Planning Policy Framework (NPPF) to maintain, restore and enhance species and habitats.
- 2.4 Seven bat species are NERC Priority Species (JNCC, 2007). These are:
- Barbastelle *Barbastella barbastellus*
 - Bechstein's *Myotis bechsteinii*
 - Noctule *Nyctalus noctula*

- Soprano Pipistrelle *Pipistrellus pygmaeus*
- Brown Long-eared *Plecotus auritus*
- Greater Horseshoe *Rhinolophus ferrumequinum*
- Lesser Horseshoe *Rhinolophus hipposideros*

2.5 Greater Horseshoe, Lesser Horseshoe, Barbastelle and Bechstein's, are afforded greater protection under European legislation, being listed under Annex II of the EC Habitats Directive which lists species whose conservation requires the designation of Special Areas of Conservation (SACs).

3.0 Methodology

Personnel

- 3.1 The survey was carried out by James Godbeer BSc Hons MCIEEM, an ecologist with over 17 years' experience working as a consultant, and experienced bat surveyors. James has extensive experience of managing environmental contracts, and particular experience in surveying, assessment and mitigation for rare and protected species. He has considerable knowledge of the development and planning process including Ecological Impact Assessments, sustainable ecological design and he has completed ecology chapters of Environmental Statements. James holds a number of protected species licences including bats (all species, all counties, Class Licence Registration No. 2015-12313-CLS-CLS), and Great Crested Newts (Class Licence Registration No. 2019-44282-CLS-CLS). He has successfully obtained European Protected Species mitigation licences for a number of bat species including Lesser Horseshoe, Greater Horseshoe, Serotine, Brown Long-eared, Common Pipistrelle and Natterer's bats, for a number of roost types including maternity and hibernation sites.

Dusk Emergence Survey

- 3.2 The buildings were subject to a dusk emergence survey on the 11th July 2024. In order to adequately cover the building, three surveyor positions were used.
- 3.3 In accordance with Bat Conservation Trust – Bat Surveys for Professional Ecologists: Good Practice Guidelines. 4th Ed (Collins, 2023), dusk surveys should begin 15 minutes before sunrise and continue for 1.5 – 2 hours after sunset with the survey start time adjusted on subsequent surveys or a repeat of the survey should bats already be in flight at 15 minutes before sunset. The dusk survey therefore began 30 minutes before sunset to avoid the risk of having to repeat the survey as species such as pipistrelles will often emerge well before sunset.

Equipment

- 3.4 Surveyors were equipped with Echo Meter Touch 2 Pro bat detectors. Registrations are recorded on the devices and notes made on species recorded, behaviour, time of registration, location and direction of flight where possible, including incidental observations from surrounding habitats. Canon XA60 video cameras used in infrared mode were used to assist with the survey.

Assessment

- 3.5 The surveyed building has been evaluated to assess which of the following categories it falls into, if any (Mitchell-Jones, 2004 & Collins, 2023):
- **Transitional roost** (April-September/October) - On waking from hibernation or in the period prior to hibernation, bats search for roosts in which they stay for only a few days or on some occasions several weeks. These transitional roosts can be occupied by a few individuals or occasionally small groups. The transitional roosts used prior to hibernation are generally cool and thus may allow bats to reduce their energy requirements before going into hibernation.
 - **Maternity roost** (May-August) - Breeding females gather together around the beginning of May to form nursery colonies. During this period gestation begins with births typically

occurring between June and July. The females and their young remain within the maternity roost until the young are weaned and independent (late July-August). These roosts tend to break up between August and September. Adult males are rarely found within these colonies. However, the adult males of long-eared bats, Daubenton's, Natterer's, and horseshoe bats can be found roosting within maternity colonies with their numbers increasing throughout the active season.

- **Satellite roost** (May-August) - Breeding females may have alternative roost sites in close proximity to the main nursery colony. These are referred to as 'satellite roosts'. The numbers of bats using these roosts can vary greatly, from a few individuals, to small groups.
- **Mating roost** (September-November) - All British bats are polygynous i.e. males mate with several females. Mating generally takes place from late summer and can continue through the winter. A number of different mating strategies are used by bats, though males of some species establish mating roosts, whereby they defend territory and display/call to females to mate.
- **Hibernation roost** (October-March) - Depending on the weather and food availability, bats tend to move to hibernation sites from October. Hibernation roosts can vary greatly in terms of the number of individuals and the diversity of species that occupy them. However, they tend to have a constant cool temperature and high humidity, which allows the bats to use less energy regulating their temperature. Bats will wake occasionally during hibernation to drink and feed.
- **Night roost** (March-November) - Bats may use roosts other than traditional day roosting sites to rest in during the night. These roosts vary in their conservation significance. Night roosts may be used by a single individual on occasion or they could be used regularly by the whole colony. Studies have shown that night roosts may be of particular importance to some species i.e. the Lesser Horseshoe, providing key resting places within core foraging areas.
- **Day roost** (March-November) - These roosts are used during the day to rest in. Males of most British species spend the summer roosting alone or in small groups with other males in such roosts. Bats may regularly use a number of day roosts, switching between them on a daily basis, though conversely they may occupy the same roosting site for several weeks.
- **Feeding roost** (May-November) - These roosts can be occupied by a single animal or a few individuals throughout the active season. They vary in their significance as they may be used by the whole colony or just a few individuals to feed, to shelter from the weather or to rest temporarily. Feeding roosts are often used by long-eared and horseshoe species.
- **Other considerations, Swarming sites** - Swarming takes place between August and November, whereby large numbers of bats from several species gather, generally around caves and mines. They are often dominated by the *Myotis* species and appear to be important mating sites with some bats travelling several kilometres to reach these areas. A proportion of the bats that travel to these sites will remain to hibernate.

Limitations

- 3.6 There were no limitations to carrying out the dusk emergence survey in accordance with the Bat Conservation Trust – Bat Surveys for Professional Ecologists: Good Practice Guidelines. 4th Ed (Collins, 2023).
- 3.7 The surveys only provide a 'snapshot' of the bat activity associated with the buildings and immediate surroundings and although the absence of a maternity roost can be determined with a high level of confidence as well as other roost types at the time of the surveys, it is possible that bats could roost at other times of the year.

4.0 Results

Dusk Emergence Survey

- 4.1 No bats were recorded emerging from, or entering the building during the survey.
- 4.2 Activity by the following species was recorded over the course of the survey (see Plan 1 for the relative distribution of activity).

Table 1: Survey results for 11th July 2024 dusk survey.

Species	Times	No. Passes	Comments
Common Pipistrelle	21:57 – 22:55	9	Individual pipistrelles recorded briefly foraging near trees to the south and over the carpark to the northeast and west.

Table 2. Weather conditions and sunset/sunrise times.

Date	Temperature (°C)	Sunset/Sunrise Times	Wind	Cloud Cover (%)
11/07/24 (dusk)	15 -13	21:25	Light	100

5.0 Evaluation

Dusk Emergence Survey

- 5.1 No bats were recorded emerging from, or entering, the building during the survey and it is therefore concluded that roosting bats were absent from the buildings at the time of the survey. The presence of a maternity roost can be ruled out with a high level of confidence and although individual bats could roost at any time, the present surveys indicate general absence.
- 5.2 One confirmed species of bat was recorded: Common Pipistrelle.
- 5.3 Occasional foraging passes by one or more Common Pipistrelles were recorded over the course of the survey, the first recorded at 32 minutes after sunset. Pipistrelle bats are the most common species of bat in the UK with widespread distributions and roosting in a wide range of habitats and building types (BCT, 2010).

Site Status Assessment

- 5.4 No roosts have been identified within the building on site. It is not possible to determine the value of the site for foraging and commuting bats based on a single dusk emergence survey. However, the building is located in urban area, surrounded predominantly by hard standing with only small pockets of colonising vegetation and therefore any foraging activity is likely to be consistently limited to individual or low numbers of more light tolerant species such as pipistrelles.

6.0 Impacts and Recommendations

Impacts

- 6.1 The proposal to redevelop the site includes the conversion of the current building to provide two new dwellings, which require significant conversion and refurbishment works.
- 6.2 In the absence of mitigation, the following impacts and potential impacts with regard to bats have been identified:
- Loss of features suitable for roosting bats; no roosts have been identified within the buildings during the surveys. No impact.
 - Potential for disturbing bats that could roost at other times of the year or those that could begin roosting at any time. Likely negligible impact.
 - Temporary/permanent disruption of small areas of bat foraging habitat on site through changes in building layout and increase/changes in external lighting but mitigation possible.

Further Surveys

- 6.3 No further surveys are required at this time. Sufficient surveys have been carried out to satisfy the recommended survey effort and repeat surveys would not be required unless the proposals are delayed by one year or more.

Legal Compliance

- 6.4 The Wildlife and Countryside Act 1981 as amended by The CROW Act 2000 and The Conservation of Habitats and Species Regulations 2010 makes it illegal to recklessly damage, destroy or obstruct access to any place that a wild bat uses for shelter or protection, whether the bat is occupying the shelter at the time or not.
- 6.5 European Protected Species (EPS) Licences to permit the above for the purposes of development must be obtained from Natural England. To gain a licence the scheme must have been issued with detailed planning permission and must not result in a loss of conservation status of the species concerned. Based on the survey evidence presented in this report it has been concluded that roosting bats are likely to be absent from the building and an **EPS Licence will not be required** to permit the works.

Mitigation

Timing of Works

- 6.6 There are no restrictions on the timing of works with respect to bats.

Care and Vigilance During Works

- 6.7 It should be noted although there was nothing to indicate the presence of roosts within the buildings on site, it is possible that crevice-dwelling bats e.g. Common Pipistrelle, could be present at other times or could begin roosting in crevices at any time, although any potential is

low. As a precaution, the contractor(s) should therefore be advised to carry out all work with care and vigilance for bats.

6.8 During the works the contractor should be advised to adhere to the following procedures in the unlikely event bats are found during works:

- If the roost is still in the structure and bats are not injured, stop work and contact a licensed ecologist. If help is not available, allow bats to fly out of harm's way.
- If material containing a roost has been removed, the roost is not exposed and the bats are not injured, temporarily seal and isolate the roost, stop work and seek advice from a licensed ecologist. If advice is not readily available, re-open it and allow bats to relocate of their own accord.
- If the roost has been exposed, and especially if bats have been injured, stop work, collect bats in a secure box or bag (using a glove) and contact a licensed ecologist

Habitat Creation

6.9 No roosting bats were recorded using the building during the survey but Common Pipistrelle activity was recorded near to the site. As no bats were found to be roosting within the building, no specific recommendation is made for new roosting features but the new development provides an opportunity to enhance the site for roosting bats by incorporating roosts into the new buildings. There are many ways that the building could be enhanced for crevice-dwelling bat species (virtually all UK bat species) without inconveniencing prospective occupants or requiring any significant design considerations. Bat tubes, panels, shelters or boxes can be installed in suitable locations around the site. A variety of aspects increases the likelihood of colonisation.

Lighting

6.10 Pipistrelle bats are generally light tolerant; but nevertheless, it is recommended that external lighting be kept to minimum in line with best practice to minimise skyglow. Measures should include the use of lighting only where absolutely necessary utilising highly directional warm white LED lighting, an example being down spots at 2.5 m high using warm white (2700 K) 8W LED lamps, 550 lumens, 35 degree beam angle. These could be individually activated by PIR sensors on a 1 minute cut off to further reduce their impacts. These will assist in lighting only the areas where lighting is required and minimising light spill either directly or through reflected light.

7.0 References

Bat Conservation Trust (2010). *Common Pipistrelle: Species information leaflet*. Bat Conservation Trust. [Online]. Available at: <https://www.bats.org.uk/about-bats/what-are-bats/uk-bats> [accessed on 21st July 2024].

Collins (2023). *Bat Surveys for Professional Ecologists: Good Practice Guidelines*. 4th Ed. The Bat Conservation Trust: London.

The Conservation of Habitats and Species Regulations 2010, SI 2010/490

The Conservation (Natural Habitats, &c.) (Amendment) Regulations 2007, SI 2007/1843, London: HMSO.

Countryside and Rights of Way Act 2000, (c.37), London: HMSO.

Defra (2011) *Biodiversity 2020: A strategy for England's wildlife and ecosystem services*. Defra.

JNCC (2007). *UK BAP Priority Species*. [Online]. Available at: <http://jncc.defra.gov.uk/page-5717> [accessed on the 21st July 2024]

Mitchell-Jones, A.J. (2004). *The Bat Mitigation Guidelines*. English Nature: Peterborough.

Wildlife and Countryside Act 1981 (and amendments), (c.69), London: HMSO.

8.0 Plans

Plan 1 – Survey Results



Surveyor Location ●

Common Pipistrelle Activity 